

Attorney Docket No.:
Inventors:
Serial No.:
Filing Date:
Page 2



PENN-0743
Greene et al.
09/783,896
February 15, 2001

RECEIVED

JUL 08 2002

TECH CENTER 1600/2900

44 10/13
11/15
11/16/02

In the Specification:

At page 1, line 1, please replace the title with the following amended title:

--Method of Detecting Molecules Expressing Selected Epitopes via
Fluorescent Dyes--

RECEIVED

JUL 05 2002

GROUP 3600

In the Claims:

1. A method for quantifying molecules expressing a selected epitope in a sample comprising:

(a) immobilizing a molecule expressing a selected epitope in a sample to a selected surface;

b¹
SUB
DI
(b) contacting the surface with an epitope detector so that the epitope detector binds to immobilized molecules on the surface, said epitope detector comprising an oligonucleotide attached to a monoclonal antibody for the selected epitope, a single chain Fv for the epitope or a constrained epitope specific CDR;

(c) amplifying the oligonucleotide of said epitope detector;

(d) contacting the amplified oligonucleotide with a fluorescent dye which binds to RNA directly and stains the

Attorney Docket No.: PENN-0743
Inventors: Greene et al.
Serial No.: 09/783,896
Filing Date: February 15, 2001
Page 3

oligonucleotide; and

(e) measuring fluorescence emitted from the stained oligonucleotide which is indicative of epitope detector bound to the surface and molecules expressing the selected epitope in the sample.

11. (amended) A method for detecting molecules expressing a selected epitope in a sample comprising:

(a) immobilizing a molecule expressing a selected epitope in a sample to a selected surface;

(b) contacting the surface with an epitope detector so that the epitope detector binds to immobilized molecules on the surface, said epitope detector comprising an oligonucleotide attached to a monoclonal antibody for the selected epitope, a single chain Fv for the epitope or a constrained epitope specific CDR;

(c) amplifying the oligonucleotide of said epitope detector;

(d) adding the amplified oligonucleotide of said epitope detector from step (c) to a reverse transcriptase based reaction or a replicase based reaction to increase sensitivity;

(e) detecting the amplified oligonucleotide of said epitope detector from step (c).